

# Waraaqda Tixraaca

Gacaliye Bixiye,

Bukaankan \_\_\_\_\_ ayaa laga yaabaa inuu qabo \_\_\_\_\_.

**Midkood doxycycline AMA ciprofloxacin ayaa loo doorbidaa dawada koowaad ee antibayotiga ee kahortagga marka cudurku jidhka soo galo kadib.** Caafimaadka dadweynaha ayaa qiimeeyay kahortagga marka cudurku jidhka soo galo ee bukaankan oo waxa loo baahanahay caawimadaada dawo-qoridda ama maaraynta dawada kahortagga ee k u habboon.

Taariikhda \_\_\_\_\_ bukaankani waxa uu holay dawada kahortagga ee markii cudurku jidhka soogalay kadib oo ah:

(taariikh)

Doxycycline 100mg po  
 BID x 10 maalmood

Ciprofloxacin 500mg po  
 BID x 10 maalmood

Wax antibiyootig ah  
 lama siin

| QAYBTA 1 | Bukaankani XASAASIYAD ayuu ku leeyahay doxy IYO                        | Kaliya Isticmaal Caafimaad   |
|----------|--|--|
|          | <input type="checkbox"/> Sidoo kale XASAASIYAD ayuu ku leeyahay cipro. | Eeg Shaxda 1, uqor antibayotig kale oo la qaato, sii bixin 10 maalmood ah  |
|          | <input type="checkbox"/> Waxa uu qaataa tizanidine (Zanaflex).         | Cipro waa mamnuuc iyadoo ay sabab u tahay inay si daran u dhexgasho dawooyinka. Yaree ama jooji tizanidine muddada lagu gudo jiro kahortagga si loo yareeyo khatarta AMA eeg Shaxda 1, uqor antibayotig kale oo la qaato, sii bixin 10 maalmood ah.    |
|          | <input type="checkbox"/> Wuxuu qabaa myasthenia gravis.                | Cipro wuxuu sii xumayn karaa murqaha daciifka ah (digniinta feerka - ka-fogaanta ayaa miyir leh). Eeg Shaxda 1, uqor antibayotig kale oo la qaato, sii bixin 10 maalmood ah AMA u qor cipro (500 mg po BID x 10 maalmood) oo si taxaddar leh ula soco. |
|          | <input type="checkbox"/> Wuxuu qabaa cudurka kalida.                   | Eeg Shaxda 2, xisaabi qiyaaska kalida ee cipro, sii 10 maalmood bixinta.   |
|          | <input type="checkbox"/> Wuxuu qabaa taariikh suuxdin/neerfo.          | Cipro waxay kordhisaa khatarta suuxdinta. Uqor cipro (500 mg po BID x 10 maalmood) oo la soco AMA Eeg Shaxda 1, uqor antibayotig kale oo la qaato, sii bixin 10 maalmood ah.   |
|          | <input type="checkbox"/> Ma awoodo inuu liqo kiniinada.                | Uqor cipro joojinta afka oo ku salaysan 15 mg/kg/qiyaasta (max 500 mg/qiyaas) 12 saacadood oo kasta, sii bixinta 10 maalmood.  |
|          | <input type="checkbox"/> Miisaankiisu wuu ka yaryahay 67 bound.        | Xisaabi qiyaasta cipro ee ku habboon iyadoo lagu saleeyay 15 mg/kg/qiista (max 500 mg/dose) 12 saacadood oo kasta, sii bixinta 10 maalmood.  |

| QAYBTA 2 | Bukaankani waxa ay leedahay UUR OO   | Kaliya Isticmaal Caafimaad   |
|----------|--|--|
|          | <input type="checkbox"/> Oo waxa ay XASAASIYAD ku qabtaa Cipro oo waxa loo qoray Doxycycline | Doxy wuxuu u horseedi karaa ilkaha wasakh iyo in ay six un u sameysmaan ilkaha carruurta hooyooyinkood ay qaatay markay uurka lahayd. Khatartu waxay ku xirantahay goorta la siinayo iyo mudada. Sii wad doxy ama eeg Eeg Shaxda 1, uqor antibayotig kale oo la qaato, sii bixin 10 maalmood ah. |
|          | <input type="checkbox"/> Waxay qaadataa tizanidine (Zanaflex).                               | Cipro waa mamnuuc iyadoo ay sabab u tahay inay si daran u dhexgasho dawooyinka. Yaree ama jooji tizanidine muddada lagu gudo jiro kahortagga si loo yareeyo khatarta AMA eeg Shaxda 1, uqor antibayotig kale oo la qaato, sii bixin 10 maalmood ah.  |
|          | <input type="checkbox"/> Waxay qabtaa myasthenia gravis.                                     | Cipro wuxuu sii xumayn karaa murqaha daciifka ah (digniinta feerka - ka-fogaanta ayaa miyir leh). Eeg Shaxda 1, uqor antibayotig kale oo la qaato, sii bixin 10 maalmood ah AMA u qor cipro (500 mg po BID x 10 maalmood) oo si taxaddar leh ula soco.   |
|          | <input type="checkbox"/> Waxay qabtaa cudur kalida ah.                                       | See Eeg Shaxda 2, xisaabi qiyaaska kalida ee cipro, sii 10 maalmood bixinta.   |
|          | <input type="checkbox"/> Waxay qabtaa taariikh suuxdin/neerfo.                               | Cipro waxay kordhisaa khatarta suuxdinta. Uqor cipro (500 mg po BID x 10 maalmood) oo la soco AMA Eeg Shaxda 1, uqor antibayotig kale oo la qaato, sii bixin 10 maalmood ah.   |
|          | <input type="checkbox"/> Ma awoodo inay liqdo kiniinada.                                     | Uqor cipro joojinta afka oo ku salaysan 15 mg/kg/qiyaasta (max 500 mg/qiyaas) 12 saacadood oo kasta, sii bixinta 10 maalmood.  |

Wixii su'aalo ah fadla la xidhiidh khadka waaxda caafimaadka:

—

# Waraqda Tixraaca

## KALIYA ISTICMAAL CAAFIMAAD

Qaybaha haray ee warqaddan gudbinta waxay ka kooban yihiin macluumaad farsamo oo loogu talagalay bixiyahaaga caafimaad oo lama turjumin.